bs-3320R

[Primary Antibody]

phospho-PDGFRA (Tyr849) / PDGFRB (Tyr857) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GenelD: 5156 **SWISS:** P16234

Target: PDGFRA (Tyr849) / PDGFRB (Tyr857)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

PDGF Receptor alpha around the phosphorylation site of Tyr849:

SN(p-Y)VS.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a cell surface tyrosine kinase receptor for

members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. Studies suggest that this gene plays a role in organ development, wound healing, and tumor progression. Mutations in this gene have been associated with idiopathic hypereosinophilic syndrome, somatic and familial gastrointestinal stromal tumors, and a variety of other cancers.

[provided by RefSeq, Mar 2012].

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500)

Reactivity: Human (predicted: Mouse,

Rat, Sheep, Cow, Chicken,

Dog, Horse)


Predicted MW.: 117 kDa

Subcellular Location: Cell membrane

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PGFRA) Polyclonal Antibody, Unconjugated (bs-3320R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell:SH-SY5Y cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Antibody incubation with (Phospho-PDGFRA (Tyr849) / PDGFRB (Tyr857)) polyclonal Antibody, Unconjugated (bs-3320R) 1:100, 90 minutes at 37°C; followed by a FITC conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

- SELECTED CITATIONS -

- [IF=8.8] Wang Haibo. et al. PDZK1 confers sensitivity to sunitinib in clear cell renal cell carcinoma by suppressing the PDGFR-β pathway. BRIT J CANCER. 2024 May;:1-14 IHC; Mouse, Human. 38822145
- [IF=2.886] Yang Q et al. Anlotinib Suppresses Colorectal Cancer Proliferation and Angiogenesis via Inhibition of AKT/ERK Signaling Cascade. Cancer Manag Res. 2020 Jun 24;12:4937-4948. WB; Mouse. 32606981