bs-0295G-SA

[Secondary Antibodies]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Reactivity: Rabbit

Goat Anti-Rabbit IgG H&L, SA conjugated

- DATASHEET -

Host: Goat Isotype: IgG

Clonality: Polyclonal

Target: Goat Anti-Rabbit IgG H&L **Purification:** affinity purified by Protein G


Concentration: 1.0 mg/ml

Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50%

glycerol.

Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

VALIDATION IMAGES -



25 ug total protein per lane of various lysates (see on figure) probed with Rabbit IgG H&L polyclonal antibody, unconjugated (bs-0295G) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

- SELECTED CITATIONS -

- [IF=3.391] Hui Yang. et al. A combined proteomic and metabolomic analyses of the priming phase during rat liver regeneration. Arch Biochem Biophys. 2020 Oct;693:108567 WB; Rat. 32898568
- [IF=3.4] Jianfei Wu. et al. A potential strategy for improving offspring behavior in maternal immune activation: Amantadine-mediated suppression of neuroinflammation. Animal Models and Experimental Medicine. 2025 Jul;: WB; Mouse. 40697163