

bs-0295G-SA

[Secondary Antibodies]

Goat Anti-Rabbit IgG H&L, SA conjugated

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Goat	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
Target: Goat Anti-Rabbit IgG H&L		IHC-F (1:100-500)
Purification: affinity purified by Protein G		IF (1:100-500)
Concentration: 1.0 mg/ml		ICC/IF (1:100-500)
Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		ELISA (1:500-1000)
		Reactivity: Rabbit

— VALIDATION IMAGES —



25 ug total protein per lane of various lysates
(see on figure) probed with Rabbit IgG H&L
polyclonal antibody, unconjugated (bs-0295G) at
1:1000 dilution and 4°C overnight incubation.
Followed by conjugated secondary antibody
incubation at r.t. for 60 min.

— SELECTED CITATIONS —

- **[IF=3.391]** Hui Yang. et al. A combined proteomic and metabolomic analyses of the priming phase during rat liver regeneration. Arch Biochem Biophys. 2020 Oct;693:108567 WB ;Rat. 32898568