

bs-11294R**[Primary Antibody]****HOXA6 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit Clonality: Polyclonal GeneID: 3203 Target: HOXA6 Immunogen: KLH conjugated synthetic peptide derived from human HOXA6/HOX1B: 171-233/233. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The Hox homeobox genes encode proteins that are transcriptional regulators with an established role in embryonic development. HoxA6 (homeobox A6), also known as HOX1B, is a 233 amino acid protein that localizes to the nucleus. Expressed during embryonic development, HoxA6 functions as a sequence-specific DNA-binding transcription factor that is part of a regulatory mechanism that provides cells with positional identities during development. Via its ability to bind DNA, HoxA6 plays an important role in the regulation of gene expression, as well as morphogenesis and differentiation. The gene encoding HoxA6 maps to human chromosome 7, which houses over 1,000 genes and comprises nearly 5% of the human genome. Defects in some of the genes localized to chromosome 7 have been linked to Osteogenesis imperfecta, Williams-Beuren syndrome, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome.	Isotype: IgG SWISS: P31267 Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Pig, Sheep, Dog, Horse) Predicted MW.: 23 kDa
--	--

— VALIDATION IMAGES —

Sample: K562(Human) Cell Lysate at 30 ug
Primary: Anti-HOXA6 (bs-11294R) at 1/500
dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 23 kD Observed band size: 26 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HOXA6) Polyclonal Antibody, Unconjugated (bs-11294R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HOXA6) Polyclonal Antibody, Unconjugated (bs-11294R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=5.145]** Deng J et al. Oxaliplatin - induced neuropathic pain involves HOXA6 via a TET1 - dependent demethylation of the SOX10 promoter. Int J Cancer. 2020 Nov 1;147(9):2503-2514. IHC ;Rat. 32428246
- **[IF=3.269]** Weifeng Wang. et al. Down-regulated long non-coding RNA LHFPL3 antisense RNA 1 inhibits the

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

radiotherapy resistance of nasopharyngeal carcinoma via modulating microRNA-143-5p/homeobox A6 axis.
Bioengineered. 2022;13(3):5421-5433 WB ;Human. 35176945