

bs-11295R**[Primary Antibody]****HOXA7 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

| | |
|--|---|
| Host: Rabbit Clonality: Polyclonal GeneID: 3204 Target: HOXA7 Immunogen: KLH conjugated synthetic peptide derived from human HOXA7: 112-230/230. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: HOX genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. The homeobox gene Hoxa-1 is transcriptionally regulated by retinoic acid (RA) and encodes a transcription factor, which has been shown to play important roles in cell differentiation and embryogenesis. Hoxa-1 is also expressed in cancers, such as mammary tumors, though it is not expressed in normal gland or in precancerous mammary tissues. At embryonic stages, Hoxa-2 is expressed in the mesenchyme and epithelial cells of palate, however its expression is restricted to the tips of the growing palatal shelves. Hoxa-2 protein is predominantly expressed in the nuclei of cells in the ventral mantle region of the developing embryo. In the developing and adult mouse spinal cord, Hoxa-2 protein may contribute to dorsal-ventral patterning and/or to the specification of neuronal phenotype. Hoxa-7 functions as a potent transcriptional repressor and its action as such requires several domains, including both activator and repressor regions. Hoxa-7 is expressed in the fetal liver, lung, skeletal muscle, kidney, pancreas and placenta. | Isotype: IgG SWISS: P31268 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse) Predicted MW.: 25 kDa Subcellular Location: Nucleus |
|--|---|

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Human liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HOXA7) Polyclonal Antibody, Unconjugated (bs-11295R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-HOXA7 Polyclonal Antibody, Unconjugated(bs-11295R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining