

**bs-10072R****[ Primary Antibody ]****VEGFB Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 7423 <b>Target:</b> VEGFB <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human VEGFB: 101-207/207. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> The onset of angiogenesis is believed to be an early event in tumorigenesis and may facilitate tumor progression and metastasis. Several growth factors with angiogenic activity have been described and include fibroblast growth factor (FGF), platelet derived growth factor (PDGF) and vascular endothelial growth factors (VEGFs). The VEGF protein family is comprised of VEGF, VEGF-B, VEGF-C and VEGF-D, all of which may exhibit angiogenic function in vivo. VEGF-B, which exists as two alternatively spliced isoforms known as VEGF-B167 and VEGF-B186, is abundantly expressed in heart and skeletal muscle and is frequently co-expressed with VEGF. VEGF-C binds to and specifically activates Flt-4 and Flk-1.	<b>Isotype:</b> IgG <b>SWISS:</b> P49765	<b>Applications:</b> Flow-Cyt (1µg/Test) <b>Reactivity:</b> Human (predicted: Mouse, Rat, Pig, Cow, Dog, GuineaPig) <b>Predicted MW.:</b> 22 kDa <b>Subcellular Location:</b> Secreted ,Extracellular matrix
---	---	---

**— VALIDATION IMAGES —**

Blank control (blue line): A549 (fixed with 2% paraformaldehyde (10 min) , then permeabilized with 90% ice-cold methanol for 30 min on ice).  
 Primary Antibody (green line): Rabbit Anti-VEGFB antibody (bs-10072R),Dilution: 1µg/10<sup>6</sup> cells;  
 Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC,Dilution: 1µg /test.