bs-5117R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Hsc70 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 3312 **SWISS:** P11142

Target: Hsc70

Immunogen: KLH conjugated synthetic peptide derived from human HSP71:

501-600/646.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a member of the heat shock protein 70 family, which contains both heat-inducible and constitutively expressed members. This protein belongs to the latter group, which are also referred to as heat-shock cognate proteins. It functions as a chaperone, and binds to nascent polypeptides to facilitate correct folding. It also functions as an ATPase in the disassembly of clathrin-coated vesicles during transport of membrane components through the cell. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1ug/test) ICC/IF (1:100-500)

Reactivity: Human, Mouse, Rat

(predicted: Rabbit, Pig,


Cow, Chicken)

Predicted 71 kDa MW.:


Subcellular Cell membrane, Cytoplasm

Location: , Nucleus

VALIDATION IMAGES




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Hsc70) Polyclonal Antibody, Unconjugated (bs-5117R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.




Blank control:U-2OS, Primary Antibody (green line): Rabbit Anti-Hsc70 antibody (bs-5117R) Dilution: 1µg /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Hsc70) Polyclonal Antibody, Unconjugated (bs-5117R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.



Blank control (black line) :Hela, Primary Antibody (green line): Rabbit Anti-Hsc70 antibody (bs-5117R) Dilution:1ug/Test; Secondary Antibody (white blue line): Goat



NIH/3T3 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Hsc70) polyclonal Antibody, Unconjugated (bs-5117R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

Antibody: Goat anti-rabbit IgG-PE Dilution: $1\mu g$ /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

anti-rabbit IgG-AF488 Dilution: 0.5ug/Test.
Isotype control (orange line): Normal Rabbit
IgG Protocol The cells were fixed with 4% PFA
(10min at room temperature) and then
permeabilized with 90% ice-cold methanol for
20 min at -20°C, The cells were then incubated in
5%BSA to block non-specific protein-protein
interactions for 30 min at room temperature
.Cells stained with Primary Antibody for 30 min
at room temperature. The secondary antibody
used for 40 min at room temperature.
Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

• [IF=3.1] Almiñana, Carmen, et al. "Oviduct extracellular vesicles protein content and their role during oviduct-embryo cross-talk." Reproduction 154.3 (2017): 153-168. WB ;Bovine. 28630101