

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## Myosin VIIa Rabbit pAb

Catalog Number: bs-7761R

Target Protein: Myosin VIIa

Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: Flow-Cyt (2ug/Test)

Reactivity: Human, Mouse (predicted:Rat, Pig, Cow, Chicken, Dog, Horse)

Predicted MW: 244 kDa Entrez Gene: 4647 Swiss Prot: Q13402

Source: KLH conjugated synthetic peptide derived from human Myosin VIIa: 851-950/2215.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins

serve in intracellular movements. Their highly divergent tails are presumed to bind to membranous compartments, which would be moved relative to actin filaments. In retina, myosin VIIa may play a role in trafficking of ribbon-synaptic vesicle complexes and renewal of the outer photoreceptors disks. In inner ear, it may maintain the rigidity of stereocilia

during the dynamic movements of the bundle.

## **VALIDATION IMAGES**



Blank control:Mouse kidney. Primary Antibody (green line): Rabbit Anti-Myosin VIIa antibody (bs-7761R) Dilution:  $2\mu g/10^6$  cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat antirabbit IgG-AF488 Dilution:  $1\mu g$ /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=10.7] Min Young Jeong. et al. Dexamethasone nanocrystals-embedded hydroxypropyl methylcellulose hydrogel increases cochlear	
delivery and attenuates hearing loss following intratympanic injection. CARBOHYD POLYM. 2024 Dec; $345:122546\ \text{IF}$ ; Mouse . $10.1016/\text{j.carbpol.}$ 2024.122546	