

bs-7772R**[Primary Antibody]****SEPT14 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Mouse, Rat (predicted: Human)
GeneID: 346288	SWISS: Q6ZU15	
Target: SEPT14		Predicted MW.: 50 kDa
Immunogen: KLH conjugated synthetic peptide derived from human SEPT14: 351-432/432.		Subcellular Location: Cytoplasm
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Septin 14, also known as SEPT14, is a 432 amino acid protein that belongs to the septin family. The highly conserved septin family of GTP-binding cytoskeletal proteins is implicated in membrane transport, apoptosis, cell polarity, cell cycle regulation, cytokinesis and other cellular functions. Septins polymerize into heterooligomeric protein complexes that form filaments, and can associate with cellular membranes, actin filaments and microtubules. Septin 14 has a GTPase domain followed by a C-terminal coiled-coil domain characteristic of group II septins. Septin 14 interacted with all septins except those that are members of its phylogenetic cluster, including itself. Septin 14 shares highest identity (67%) with Septin 10. Expressed in testis and more weakly in fetal liver, tonsil and thymus, the Septin 14 protein is not detected in testicular cancer or other normal or cancer cell lines. The Septin 14 gene is conserved in chimpanzee, dog, cow, mouse and rat, and maps to human chromosome 7p11.2.		

— VALIDATION IMAGES —

Sample: Testis (Mouse) Lysate at 40 ug Primary:

Anti- SEPT14 (bs-7772R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at

1/20000 dilution Predicted band size: 50 kD

Observed band size: 50 kD