

bs-11379R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Syntenin 1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 6386	SWISS: O00560	IF (1:100-500)
Target: Syntenin 1		Reactivity: Mouse (predicted: Human, Rat, Rabbit, Chicken, Dog)
Immunogen: KLH conjugated synthetic peptide derived from human Syntenin 1: 121-220/298.		
Purification: affinity purified by Protein A		Predicted MW.: 32 kDa
Concentration: 1mg/ml		Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Syntenin-1 (also known as syntenin, syndecan binding protein, melanoma differentiation-associated protein 9 or proTGF-alpha cytoplasmic domain-interacting protein 18) is a protein that binds to the cytoplasmic domains of the syndecans in yeast 2-hybrid screens and other assays. Syntenin-1 contains a tandem repeat of PDZ domains that reacts with the FYA (phe-tyr-ala) carboxy-terminal amino acid sequence of the syndecans. It may function as an adaptor that couples syndecans to cytoskeletal proteins or cytosolic downstream signal-effectors. Syntenin-1 colocalizes and interacts specifically with immature, intracellular forms of proTGF-alpha. It is a human gamma interferon responsive protein. Syntenin-1 contains PSD-95/Discs large/zO-1 (PDZ) domains and associates with the cytoplasmic tail of the IL-5Ralpha. It directly associates with the transcription factor Sox-4. The PDZ proteins PICK1, GRIP, ABP and syntenin-1 bind multiple glutamate receptor subtypes.		

— VALIDATION IMAGES —

Tissue/cell: Mouse Embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Syntenin 1 Polyclonal Antibody, Unconjugated(bs-11379R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining