

**bs-11142R****[ Primary Antibody ]****Rho B Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500)
<b>Clonality:</b> Polyclonal		<b>IHC-F</b> (1:100-500)
<b>GeneID:</b> 388	<b>SWISS:</b> P62745	<b>IF</b> (1:100-500)
<b>Target:</b> Rho B		<b>Reactivity:</b> Human (predicted: Mouse, Rat, Rabbit, Chicken, Dog)
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Rho B: 101-196/196.		
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 22 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Cell membrane ,Cytoplasm ,Nucleus
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The Rho proteins interact directly with protein kinases which are likely to serve as downstream effector targets of the activated GTPase. The Rho family of GTPases are involved in the regulation of the actin skeleton. Activation of Rho to the GTP bound state promotes formation of stress fibers and is required for the formation and maintenance of focal adhesions. Pathways stimulated by cytokines, hormones and various forms of stress utilize p21 proteins of the Rho family. Possible mediators for these responses include protein kinase such as p120PKN, and p150 ROKa/Rho-A kinase, myosin light chain (MLC) phosphatase and PIP 5-kinase.		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Human pancreatic cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Rho B) Polyclonal Antibody, Unconjugated (bs-11142R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human pancreatic cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Rho B) Polyclonal Antibody, Unconjugated (bs-11142R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

**— SELECTED CITATIONS —**

- **[IF=2.1]** Jun Li. et al. WJ-MSCs intervention may relieve intrauterine adhesions in female rats via TGF-β1-mediated Rho/ROCK signaling inhibition. Mol Med Rep. 2021 Jan;23(1):1-1 IHC ;Rat. 33179074