bs-19129R

- DATASHEET -

[Primary Antibody]

ZFP64 Rabbit pAb

ANTIB www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

Reactivity: Mouse, Rat (predicted: Human, Pig, Cow, Dog, Horse)

Predicted MW.: 75 kDa

Subcellular Location: Nucleus

Host: Rabbit Clonality: Polyclonal

SWISS: Q9NPA5

Isotype: IgG

GenelD: 55734 Target: ZFP64

Immunogen: KLH conjugated synthetic peptide derived from human ZFP64: 391-490/681.

Purification: affinity purified by Protein A


Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated


freeze/thaw cycles.

Background: ZFP64 is a 681 amino acid homolog of the mouse Zfp64 protein and is a member of the Krüppel C2H2-type zinc-finger family. Localized to the nucleus, ZFP64 contains nine C2H2-type zinc fingers and is thought to be involved in transcriptional regulation. Four isoforms of ZFP64 exist due to alternative splicing events.

- VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Rat bladder); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP64) Polyclonal Antibody, Unconjugated (bs-19129R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP64) Polyclonal Antibody, Unconjugated (bs-19129R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.