bs-19135R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

ZFP92 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 139735 **SWISS:** A6NM28

Target: ZFP92

Immunogen: KLH conjugated synthetic peptide derived from human ZFP92:

301-400/416.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: ZFP92 belongs to the krueppel C2H2-type zinc-finger protein

family. It contains eight C2H2-type zinc fingers and one KRAB domain. ZFP92 may be involved in transcriptional regulation.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500)


Reactivity: Human, Mouse

(predicted: Rabbit)

Predicted MW.: 406 kDa

Subcellular Nucleus

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP92) Polyclonal Antibody, Unconjugated (bs-19135R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP92) Polyclonal Antibody, Unconjugated (bs-19135R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.