

---

## ZFP92 Rabbit pAb

Catalog Number: bs-19135R

Target Protein: ZFP92

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse (predicted:Rabbit)

Predicted MW: 406 kDa

Subcellular Nucleus

Locations:

Entrez Gene: 139735

Swiss Prot: A6NM28

Source: KLH conjugated synthetic peptide derived from human ZFP92: 301-400/416.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: ZFP92 belongs to the krueppel C2H2-type zinc-finger protein family. It contains eight C2H2-type zinc fingers and one KRAB domain. ZFP92 may be involved in transcriptional regulation.

### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP92) Polyclonal Antibody, Unconjugated (bs-19135R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP92) Polyclonal Antibody, Unconjugated (bs-19135R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.