

bs-14197R**[Primary Antibody]****DARS Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse (predicted: Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse) Predicted MW.: 57 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 1615	SWISS: P14868	
Target: DARS		
Immunogen: KLH conjugated synthetic peptide derived from human Cell proliferation inducing protein 40: 401-501/501.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Aspartyl-tRNA synthetase (DARS) is part of a multienzyme complex of aminoacyl-tRNA synthetases. Aspartyl-tRNA synthetase charges its cognate tRNA with aspartate during protein biosynthesis. [provided by RefSeq, Jul 2008]		

— VALIDATION IMAGES —

Sample: Liver (Mouse) Lysate at 40 ug Primary:
Anti- DARS (bs-14197R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 57 kD
Observed band size: 57 kD



Paraformaldehyde-fixed, paraffin embedded
(Human liver cancer); Antigen retrieval by boiling
in sodium citrate buffer (pH6.0) for 15min; Block
endogenous peroxidase by 3% hydrogen
peroxide for 20 minutes; Blocking buffer (normal
goat serum) at 37°C for 30min; Antibody
incubation with (DARS) Polyclonal Antibody,
Unconjugated (bs-14197R) at 1:400 overnight at
4°C, followed by operating according to SP
Kit(Rabbit) (sp-0023) instructions and DAB
staining.

— SELECTED CITATIONS —

- **[IF=3.8]** Xiong Hao. et al. DARS expression in BCR/ABL1-negative myeloproliferative neoplasms and its association with the immune microenvironment. SCI REP-UK. 2024 Jul;14(1):1-11 IHC ;Human. 39030308