

bs-12503R**[Primary Antibody]****APPBP1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit Clonality: Polyclonal GeneID: 8883 Target: APPBP1 Immunogen: KLH conjugated synthetic peptide derived from human APPBP1: 401-500/534. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: APPBP1 is a member of the ubiquitin-activating E1 family. In fetal tissues APPBP1 is widely expressed and in adult tissues it is expressed throughout the brain. APPBP1 is a cell membrane associated protein and functions as the regulatory subunit in a heterodimer with UBA3. The APPBP1/UBA3 complex binds to and activates NEDD8, a ubiquitin-like protein involved in signal transduction, cell proliferation and development. This suggests that APPBP1 affects a variety of cellular functions. In addition, APPBP1 is essential for cell cycle progression through the S/M checkpoint. More specifically, it inhibits the entry into S phase and promotes entry into M phase.	Isotype: IgG SWISS: Q13564	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Dog, Horse) Predicted MW.: 60 kDa Subcellular Location: Cell membrane
--	---	--

— VALIDATION IMAGES —

Sample: Placenta (Mouse) Lysate at 40 ug
 Primary: Anti-APPBP1 (bs-12503R) at 1/300
 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 60 kD Observed band size: 60 kD



Paraformaldehyde-fixed, paraffin embedded (human myocardium); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (APPBP1) Polyclonal Antibody, Unconjugated (bs-12503R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (APPBP1) Polyclonal Antibody, Unconjugated (bs-12503R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=2.1]** Xiangdong Meng. et al. Naringin ameliorates memory deficits and exerts neuroprotective effects in a mouse model of Alzheimer's disease by regulating multiple metabolic pathways. Mol Med Rep. 2021 May;23(5):1-13 WB ;Mouse. 33760152