

bs-2096R**[Primary Antibody]****SLC4A4 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

| | |
|--|--|
| <p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 8671</p> <p>Target: SLC4A4</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human SLC4A4 isoform 2: 1-100/1035. < Cytoplasmic ></p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: SLC4A4 (Electrogenic sodium bicarbonate cotransporter 1) is an electrogenic sodium/bicarbonate cotransporter with a Na(+):HCO3(-) stoichiometry varying from 1:2 to 1:3. It may regulate bicarbonate influx/efflux at the basolateral membrane of cells and regulate intracellular pH. SLC4A4 interacts with carbonic anhydrase 2 and carbonic anhydrase 4 which may regulate transporter activity. There are four named isoforms produced by alternative splicing.</p> <p>This gene encodes a sodium bicarbonate cotransporter (NBC) involved in the regulation of bicarbonate secretion and absorption and intracellular pH. Mutations in this gene are associated with proximal renal tubular acidosis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008].</p> | <p>Applications: WB (1:500-2000)</p> <p>Reactivity: Human, Mouse (predicted: Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)</p> <p>Predicted MW.: 116 kDa</p> <p>Subcellular Location: Cell membrane</p> |
|--|--|

— VALIDATION IMAGES —

Sample: Kidney (Mouse) Lysate at 40 ug Primary:
Anti- SLC4A4 (bs-2096R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 116 kD
Observed band size: 116 kD

— SELECTED CITATIONS —

- **[IF=3.553]** Liu M et al. Copper promotes sheep pancreatic duct organoid growth by ATOX1-dependent MEK-ERK pathway activation. Am J Physiol Cell Physiol. 2020 Apr 1;318(4):C806-C816. IF ;sheep. 32130071
- **[IF=3.553]** Liu M et al. Copper promotes sheep pancreatic duct organoid growth by ATOX1-dependent MEK-ERK pathway activation. Am J Physiol Cell Physiol. 2020 Apr 1;318(4):C806-C816. IF ;sheep. 32130071