

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

UCP-2 Rabbit pAb

Catalog Number: bs-20751R

Target Protein: UCP-2 Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW: 34 kDa Entrez Gene: 7351 Swiss Prot: P55851

Source: KLH conjugated synthetic peptide derived from human UCP-2: 51-120/309.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: UCPs facilitate the transfer of anions from the inner to the outer mitochondrial membrane

and the return transfer of protons from the outer to the inner mitochondrial membrane.


They also reduce the mitochondrial membrane potential in mammalian cells. UCP2 gene is expressed in many tissues, with the greatest expression in skeletal muscle. UCP2 is thought

to play a role in non shivering thermogenesis, obesity and diabetes.


VALIDATION IMAGES



Sample: Hela(Human) Cell Lysate at 30 ug Primary: Anti-UCP-2 (bs-20751R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 34 kD



Sample: Skeletal muscle (Mouse) Lysate at 40 ug Primary: Anti- UCP-2 (bs-20751R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 35 kD



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UCP-2) Polyclonal Antibody, Unconjugated (bs-20751R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=3.038] Lan Yang et al. The Involvement of Mitochondrial Biogenesis in Selenium Reduced Hyperglycemia-Aggravated Cerebral Ischemia Injury. Neurochem Res. 2020 Aug;45(8):1888-1901. WB; MOUSE . 32447509

[IF=2.22] Ya Sun. et al. Expression and hormone regulation of UCP2 in goat uterus. ANIM REPROD SCI. 2022 Aug;243:107015 WB,IHC; Goat . 35689907