bs-20751R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

UCP-2 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 7351 **SWISS:** P55851

Target: UCP-2

Immunogen: KLH conjugated synthetic peptide derived from human UCP-2:

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: UCPs facilitate the transfer of anions from the inner to the outer

mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. UCP2 gene is expressed in many tissues, with the greatest expression in skeletal muscle. UCP2 is thought to play a role in non shivering

thermogenesis, obesity and diabetes.

Applications: WB (1:500-2000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)


Reactivity: Human, Mouse, Rat

(predicted: Rabbit, Pig, Sheep, Cow, Dog, Horse)


Predicted 34 kDa

Subcellular Cytoplasm Location:


VALIDATION IMAGES



Sample: Hela(Human) Cell Lysate at 30 ug Primary: Anti-UCP-2 (bs-20751R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 34 kD



Sample: Skeletal muscle (Mouse) Lysate at 40 ug Primary: Anti- UCP-2 (bs-20751R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 35 kD



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UCP-2) Polyclonal Antibody, Unconjugated (bs-20751R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

— SELECTED CITATIONS —

- [IF=5.4] Hai Liu. et al. Protective effects of Ginkgolide B on myocardial ischemia reperfusion injury: role of the GAS6/Axl signaling pathway. CHEM-BIOL INTERACT. 2025 Jun;:111607 WB; Mouse. 40541647
- [IF=3.038] Lan Yang et al. The Involvement of Mitochondrial Biogenesis in Selenium Reduced Hyperglycemia-Aggravated Cerebral Ischemia Injury. Neurochem Res. 2020 Aug;45(8):1888-1901. WB; Mouse. 32447509
- [IF=2.22] Ya Sun. et al. Expression and hormone regulation of UCP2 in goat uterus. ANIM REPROD SCI. 2022 Aug;243:107015 WB,IHC;Goat. 35689907