bsm-54216R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

TYRP1 Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Recombinant

GeneID: 7306 SWISS: P17643

Target: TYRP1

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: TRP1 is a melanosomal enzyme that belongs to the tyrosinase

family and plays an important role in the melanin biosynthetic pathway. Defects in this gene are the cause of rufous

oculocutaneous albinism and oculocutaneous albinism type III.

Applications: WB (1:500-2000)


IHC-P (1:100-200) IHC-F (1:100-200) **IF** (1:100-200) **IP** (1:20-50)

Reactivity: Human

Predicted MW.: 58 kDa

Subcellular Location: Cytoplasm

VALIDATION IMAGES



Immunohistochemical analysis of paraffinembedded human malignant melanoma tissue using anti-TRP1 antibody. The section was pretreated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (bsm-54216R, 1/400) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunohistochemical analysis of paraffinembedded human skin tissue using anti-TRP1 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (bsm-54216R, 1/100) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.