bs-10343R

## [ Primary Antibody ]

## Phosphor-PPP3CA (Ser469) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GeneID:** 5530 **SWISS:** Q08209

Target: Phosphor-PPP3CA (Ser469)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

PPP3CA around the phosphorylation site of Ser469: 421-521/521.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Calcineurin, a major soluble calmodulin binding protein in the brain, a Ca2+/calmodulin dependent serine/threonine protein phosphatase, with a relatively narrow substrate specificity. This metalloenzyme, also known as phosphatase 2B, is a heterodimer composed of a calmodulin binding, catalytic alpha subunit 61 kDa, calcineurin A) and calcium binding beta subunit (18 kDa, calcineurin B). The Ca2+ binding subunit, calcineurin B, is immunologically conserved from yeast to mammals. The presence of 2 different calcineurin B isoforms (beta 1 and beta 2) has been reported in rat testis. The catalytic subunit of calcineurin, calcineurin A, isolated from different tissues or different organisms, exhibits some immunological heterogeneity. Further, there are at least 2 isoforms of calcineurin A in bovine brain (alpha 1 and alpha 2, 61 and 59 kDa, respectively).

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Mouse (predicted: Human,


Rat, Rabbit, Cow, Chicken,

GuineaPig)

**Predicted** 57 kDa MW.:

Subcellular Nucleus

## **VALIDATION IMAGES**



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PPP3CA (Ser469)) Polyclonal Antibody, Unconjugated (bs-10343R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.