
Prostaglandin I2 Receptor Rabbit pAb

Catalog Number: bs-11307R

Target Protein: Prostaglandin I2 Receptor

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: ELISA (1:5000-10000)

Reactivity: Human (predicted:Mouse, Rat, Rabbit, Pig, Cow, Dog, Horse)

Predicted MW: 41 kDa

Entrez Gene: 5739

Swiss Prot: P43119

Source: KLH conjugated synthetic peptide derived from human Prostaglandin I2 Receptor:
101-200/386.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Cyclooxygenases metabolize arachidonate to five primary prostanoids: PGE2, PGF2?, PGI2, TXA2 and PGD2. These lipid mediators interact with specific members of G protein-coupled prostanoid receptors, designated EP, FP, IP, TP and DP, respectively. The IP Receptor binds prostacyclin, PGI2, the main pro-stanoid synthesized by vascular tissues. Upon binding to the IP Receptor, prostacyclin activates adenylate cyclase primarily through the Gas protein. The gene encoding the human IP Receptor is located on chromosome 19. It is expressed as a glycosylated and phosphorylated protein, which is abundantly expressed in vascular tissues such as aorta, lung, atrium and ventricle, as well as in kidney, thymus, spleen and neurons.