### bsm-54217R

## [ Primary Antibody ]

# Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# CSNK2B/CK II beta Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 2C9
GeneID: 1460 SWISS: P67870

Target: CSNK2B/CK II beta

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** This gene encodes the beta subunit of casein kinase II, a

ubiquitous protein kinase which regulates metabolic pathways, signal transduction, transcription, translation, and replication. The enzyme is composed of three subunits, alpha, alpha prime and beta, which form a tetrameric holoenzyme. The alpha and alpha prime subunits are catalytic, while the beta subunit serves regulatory functions. The enzyme localizes to the endoplasmic reticulum and the Golgi apparatus. [provided by RefSeq, Jul 2008].

**Applications: WB** (1:500-2000)


IHC-P (1:50-100) IHC-F (1:400-800) IF (1:100-500)

Reactivity: Human, Mouse, Rat


Predicted MW.: 25 kDa

Subcellular Nucleus


#### - VALIDATION IMAGES -



Sample: Lane 1: Mouse Cerebellum tissue lysates Lane 2: Mouse Cerebrum tissue lysates Lane 3: Mouse Testis tissue lysates Lane 4: Rat Cerebellum tissue lysates Lane 5: Rat Cerebrum tissue lysates Lane 5: Rat Cerebrum tissue lysates Lane 6: Rat Kidney tissue lysates Lane 7: Rat Testis tissue lysates Lane 8: Human Hela cell lysates Lane 9: Human Jurkat cell lysates Lane 10: Human K562 cell lysates Lane 11: Human HepG2 cell lysates Lane 12: Human MCF-7 cell lysates Lane 13: Human SH-SY5Y cell lysates Primary: Anti-CSNK2B/CK II beta (bsm-54217R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 25 kDa Observed band size: 25 kDa



Paraformaldehyde-fixed, paraffin embedded (human brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CSNK2B CK II beta) Monoclonal Antibody, Unconjugated (bsm-54217R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CSNK2B CK II beta) Monoclonal Antibody, Unconjugated (bsm-54217R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.