
GARB1 Rabbit pAb

Catalog Number: bs-3766R

Target Protein: GARB1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Cow, Chicken, Dog, Horse)

Predicted MW: 47 kDa

Entrez Gene: 2554

Swiss Prot: P14867

Source: KLH conjugated synthetic peptide derived from human GABA A Receptor beta 1: 351-456/456.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (g-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl⁻ conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R alpha 1-6, GABAA R beta 1-3, GABAA R ρ 1-3, GABAA R δ , GABAA R gamma, GABAA R delta 1 and GABAA R delta 2. The GABAB family is composed of GABAB R1 alpha and GABAB R1 beta. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2 and -3). The GABA transporters function to terminate GABA action.

VALIDATION IMAGES

Sample: brain (Mouse) Lysate at 40 ug Primary: Anti-GARB1 (bs-3766R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 47 kD Observed band size: 47 kD



Sample: Cerebrum (Rat) Lysate at 40 ug Cerebrum (Mouse) Lysate at 40 ug Cerebellum (Rat) Lysate at 40 ug
Primary: Anti- GARB1 (bs-3766R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 47 kD Observed band size: 47 kD



PRODUCT SPECIFIC PUBLICATIONS

[IF=2.7] Won Seob Kim. et al. Exploring the impact of temporal heat stress on skeletal muscle hypertrophy in bovine myocytes. J THERM BIOL. 2023 Aug;:103684 IF ; Bovine . 37625343