

bs-21033R**[Primary Antibody]****SLIRP/C14orf156 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 81892**SWISS:** Q9GZT3**Target:** SLIRP/C14orf156**Immunogen:** KLH conjugated synthetic peptide derived from human SLIRP/C14orf156: 31-100/109.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Steroid receptor RNA activator (SRA, or SRA1; MIM 603819) is a complex RNA molecule containing multiple stable stem-loop structures that functions in coactivation of nuclear receptors. SLIRP interacts with stem-loop structure-7 of SRA (STR7) and modulates nuclear receptor transactivation (Hatchell et al., 2006 [PubMed 16762838]).[supplied by OMIM, Mar 2008]**Applications:** **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Human, Mouse
(predicted: Rat, Rabbit, Pig)**Predicted
MW.:** 12 kDa**Subcellular
Location:** Nucleus**— VALIDATION IMAGES —**


Sample: Kidney (Mouse) Lysate at 40 ug Primary:
Anti- SLIRP/C14orf156 (bs-21033R) at 1/1000
dilution Secondary: IRDye800CW Goat Anti-
Rabbit IgG at 1/20000 dilution Predicted band
size: 12 kD Observed band size: 11 kD



Sample: Testis (Mouse) Lysate at 40 ug Primary:
Anti-C14orf156 (bs-21033R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 12 kD
Observed band size: 12 kD



Paraformaldehyde-fixed, paraffin embedded
(Human liver carcinoma); Antigen retrieval by
boiling in sodium citrate buffer (pH6.0) for
15min; Block endogenous peroxidase by 3%
hydrogen peroxide for 20 minutes; Blocking
buffer (normal goat serum) at 37°C for 30min;
Antibody incubation with (SLIRP/C14orf156)
Polyclonal Antibody, Unconjugated (bs-21033R)
at 1:400 overnight at 4°C, followed by operating
according to SP Kit(Rabbit) (sp-0023)
instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded
(Mouse liver); Antigen retrieval by boiling in
sodium citrate buffer (pH6.0) for 15min; Block
endogenous peroxidase by 3% hydrogen
peroxide for 20 minutes; Blocking buffer (normal
goat serum) at 37°C for 30min; Antibody
incubation with (SLIRP/C14orf156) Polyclonal
Antibody, Unconjugated (bs-21033R) at 1:400

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

overnight at 4°C, followed by operating
according to SP Kit(Rabbit) (sp-0023)
instructions and DAB staining.