

bs-4622R

[Primary Antibody]

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Mycobacterium bovis Rabbit pAb

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>Target: Mycobacterium bovis</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Antigen 85B is the most abundant protein expressed by Mycobacterium tuberculosis (about one quarter). It is a mycolyl transferase in the myc pathway and catalyses - like Ag85A and Ag85C - the transfer of the fatty acid mycolate from one trehalose monomycolate to another, resulting in trehalose dimycolate and free trehalose and helping build the cell wall.</p>	<p>Isotype: IgG</p>	<p>Applications: ELISA (1:5000-10000)</p> <p>Reactivity: (predicted: Cow, Mycobacteriumis)</p> <p>Subcellular Location: Secreted</p>
--	----------------------------	---

— SELECTED CITATIONS —

- **[IF=5.6]** Lucia F. Saad. et al. A disposable, portable electrochemical immunosensor for rapid in situ detection of bovine tuberculosis. TALANTA. 2025 Jan;281:126878 ;Bovine. 39276570
- **[IF=1.85]** Seo, Min-Goo, et al. "Mycobacterium tuberculosis Infection in a Domesticated Korean Wild Boar (Sus scrofa coreanus)." Journal of Food Protection 80.6 (2017): 1009-1014. Other ;="". 28485632
- **[IF=0.994]** Shin-ichi Nakamura. et al. Eosinophilic Leukaemia and Systemic Mycobacterium marinum Infection in an African Pygmy Hedgehog (Atelerix albiventris). J Comp Pathol. 2020 Nov;181:33 IHC ;Hedgehog. 33288148
- **[IF=0]** Di Blasio A et al. Animal tuberculosis in a free-ranging fallow deer in northwest Italy: a case of "lucky strain survival" or multi-host epidemiological system complexity? European Journal of Wildlife Research,2019 65(5). IHC ;Deer. doi:10.1007/s10344-019-1316-0