bs-11410R

[Primary Antibody]

Progestin Receptor Beta Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 85315 **SWISS:** Q8TEZ7

Target: Progestin Receptor Beta

Immunogen: KLH conjugated synthetic peptide derived from human Progestin

Receptor Beta: 251-300/354. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The steroid progesterone induces the resumption of maturation in oocytes via a nongenomic pathway through binding to a novel membrane progestin receptor (mPR). This pathway inhibits adenylyl cyclase and reduces intracellular cAMP, and also activates mitogen-activated protein kinase to effect signal transduction pathways. Five distinct groups, designated Alpha, Beta, Gamma, Delta, comprise the mPR gene family. mPR Alpha, also designated progestin and adipo Theta receptor family member VII (PAQR7), consists of an extracellular N-terminus, an intracellular C-terminus and seven transmembrane domains. mPR Alpha is expressed in ovary, testis, placenta, uterus and bladder. mPR Beta, or progestin and adipo Theta receptor family member VIII (PAQR8), consists of eight putative transmembrane regions and an intracellular Nterminus that contains a leucine-rich motif. mPR Beta is a 354 amino acid protein expressed in brain and spinal cord. Both mPR Alpha and mPR Beta may be G protein-coupled receptors and may be involved in oocyte maturation.


Applications: WB (1:500-2000)

Reactivity: Human, Mouse

Predicted 40 kDa MW.:

Subcellular Cell membrane

VALIDATION IMAGES -



25 ug total protein per lane of various lysates (see on figure) probed with Progestin Receptor Beta polyclonal antibody, unconjugated (bs-11410R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

— SELECTED CITATIONS —

- [IF=17.173] Thomas Topilko. et al. Edinger-Westphal peptidergic neurons enable maternal preparatory nesting. Neuron. 2022 Feb;: IHC; Mouse. 35123655
- [IF=4.26] Kasubuchi, Mayu, et al. "Membrane progesterone receptor beta (mPRβ/Paqr8) promotes progesteronedependent neurite outgrowth in PC12 neuronal cells via non-G protein-coupled receptor (GPCR) signaling." Scientific Reports 7 (2017). WB; Mouse. 28701790

- [IF=3.7] Thiago Almeida-Souza. et al.Sex differences in long-term fear and anxiety-like responses to deep brain stimulation in a preclinical model of PTSD.journal of psychiatric research.2025 Feb 26:184:198-209. Western Blot; Rat. 40056639
- [IF=2.299] Gao XX et al. Effects of l-arginine on endometrial estrogen receptor α/β and progesterone receptor expression in nutrient-restricted sheep. Theriogenology. 2019 Jul 25;138:137-144. IHC, IF, WB; ewe. 31352175