bs-11420R

[Primary Antibody]


NMUR1 Rabbit pAb



sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	lsotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Mouse, Rat
GenelD: 10316	SWISS: Q9HB89	
Target: NMUR1		
Immunogen: KLH conjugated synthetic peptide derived from human NMUR1: 31-80/426. < Extracellular >		Predicted MW.: 47 kDa
Purification: affinity purified by	Protein A	Subsellular
Concentration: 1mg/ml		Subcellular Location: Cell membrane
Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Neuromedin U is a neuropeptide with high activity on smooth muscle. It is widely expressed in gastrointestinal systems and central nervous system (CNS). Peripheral activities of neuromedin U include smooth muscle stimulation, ion transport alterations in the gut and the regulation of local blood flow and adrenocortical function. Neuromedin U receptors 1 and 2 (NMUR1 and NMUR2) are multi-pass membrane proteins that belong to the G-protein coupled receptor 1 family of proteins. Both NMUR1 and NMUR2 act as receptors for the neuromedin U neuropeptide. NMUR1 is detected in peripheral organs, particularly in urogenital and gastrointestinal systems, with highest levels in testis. It' s expression in CNS is low, but the protein has been detected in cerebellum, hippocampus, dorsal root ganglia and spinal cord. NMUR2 is predominantly detected in central nervous system with highest levels detected in medulla oblongata, spinal cord and thalamus. It may also be detected in testis but has low levels of expression in peripheral tissues.		edin ns in ical t2) in R2 act d. with

- VALIDATION IMAGES ------



25 ug total protein per lane of various lysates (see on figure) probed with NMUR1 polyclonal antibody, unconjugated (bs-11420R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.