
ARFGAP3 Rabbit pAb

Catalog Number: bs-12513R

Target Protein: ARFGAP3

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Mouse, Rat (predicted:Human, Sheep, Cow, Chicken, Dog)

Predicted MW: 57 kDa

Entrez Gene: 26286

Swiss Prot: Q9NP61

Source: KLH conjugated synthetic peptide derived from human ARFGAP3: 412-516/516.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: ARFGAP1 (ADP-ribosylation factor GTPase-activating protein 1), ARFGAP2 (ADP-ribosylation factor GTPase-activating protein 2) and ARFGAP3 (ADP-ribosylation factor GTPase-activating protein 3) are GTPase-activating proteins (GAP) that are associated with the Golgi apparatus and interact with ADP-ribosylation factor 1 (ARF). These proteins promote hydrolysis of ARF-bound GTP and are required for the dissociation of coat proteins from Golgi-derived membranes and vesicles. Dissociation of the coat proteins is required for the fusion of these vesicles with target compartments. The activity of ARFGAP1, ARFGAP2 and ARFGAP3 is stimulated by phosphoinositides and inhibited by phosphatidylcholine. The genes encoding ARFGAP1, ARFGAP2 and ARFGAP3 map to human chromosomes 20q13.33, 11p11.2 and 22q13.2, respectively.

VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with ARFGAP3 polyclonal antibody, unconjugated (bs-12513R) at 1:500 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.