
A4GALT Rabbit pAb

Catalog Number: bs-7588R

Target Protein: A4GALT

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: Flow-Cyt (1ug/Test)

Reactivity: Human, Mouse (predicted:Rat, Pig, Chicken, Dog)

Predicted MW: 40 kDa

Entrez Gene: 53947

Swiss Prot: Q9NPC4

Source: KLH conjugated synthetic peptide derived from human A4GALT/CD77: 291-353/353.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Necessary for the biosynthesis of the Pk antigen of blood histogroup P. Catalyzes the transfer of galactose to lactosylceramide and galactosylceramide. Necessary for the synthesis of the receptor for bacterial verotoxins. Expression of CD77, also called Gb3, sensitizes a cell to verotoxins, causing cellular injury that can lead to disease. Therefore, the complex regulation of CD77 biosynthesis and the activity of the enzymes involved, such as CD77 synthase, can be studied by compared gene expression between toxin-sensitive and insensitive tissues and cell lines. The highest tissue expression of CD77 synthase occurs in the kidney, mesenteric lymph node, spleen, and brain. Burkitt leukemia cells express very high levels of CD77 as well as CD77 synthase, and are sensitive to verotoxin induced apoptosis. These megakaryoblasts then never mature, leading to the arrest of platelet generation in the bone marrow, which may cause thrombocytopenia, a symptom associated with various hemorrhagic conditions.

VALIDATION IMAGES



Blank control(blue):mouse spleen cells (fixed with 2% paraformaldehyde (10 min), then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody: Rabbit Anti- A4GALT antibody(bs-7588R), Dilution: 1µg in 100 µL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions); Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

PRODUCT SPECIFIC PUBLICATIONS

[IF=3.4] Zhen Xiao. et al. Identification and immunological characterization of genes associated with ferroptosis in Alzheimer's disease and experimental demonstration. MOL MED REP. 2024 Sep;30(3):1-15 WB ; Human . 38963039