### bs-21140R

### [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## – DATASHEET –

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 729475 **SWISS:** Q09MP3

Target: RAD51AP2

Immunogen: KLH conjugated synthetic peptide derived from human RAD51AP2:

151-250/1159.

RAD51AP2 Rabbit pAb

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS,

pH1.4.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500)


Reactivity: Rabbit, Dog

(predicted: Human, Horse)


Predicted MW.: 134 kDa

Subcellular Cytoplasm ,Nucleus

#### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Dog testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RAD51AP2) Polyclonal Antibody, Unconjugated (bs-21140R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rabbit small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RAD51AP2) Polyclonal Antibody, Unconjugated (bs-21140R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.