

CNKS2/KSR2 Rabbit pAb

Catalog Number: bs-8029R

Target Protein: CNKS2/KSR2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Cow, Chicken, Dog)

Predicted MW: 114 kDa

Entrez Gene: 22866

Swiss Prot: Q8WXI2

Source: KLH conjugated synthetic peptide derived from human CNKS2 corresponding to amino acids 863-508.: 481-580/1034.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: CNKS2 may function as an adapter protein or regulator of Ras signaling pathways.


VALIDATION IMAGES



Sample: U251(Human) Cell Lysate at 30 ug Primary: Anti-CNKS2/KSR2 (bs-8029R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 114 kD Observed band size: 114 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CNKS2/KSR2) Polyclonal Antibody, Unconjugated (bs-8029R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CNKSR2/KSR2) Polyclonal Antibody, Unconjugated (bs-8029R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.