bs-11331R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

TBR2 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 8320 **SWISS:** 095936

Target: TBR2

Immunogen: KLH conjugated synthetic peptide derived from human TBR2:

301-400/686.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: EOMES (eomesodermin homolog), also known as TBR2 (T-box-

brain2), is the 686 amino acid human homolog of the mouse Eomes protein that contains one T-box DNA binding domain. Genes that contain T-box domains encode proteins that function as transcription factors and are often involved in the regulation of various developmental events. Localized to the nucleus and expressed in the developing brain, EOMES is thought to be involved in neuronal migration and division and may play a role in trophoblast development and gastrulation. Silencing of the EOMES gene can cause mutated or arrested development and may lead to microcephaly disorders which are characterized by reduced head

circumference and a malformed brain.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test)

Reactivity: Human, Rat

(predicted: Mouse, Sheep, Cow, Chicken, Horse)

Predicted 72


MW.: ^{73 kDa}

Subcellular Nucleus **Location:**

VALIDATION IMAGES



Tissue/cell: rat brain tissue; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-TBR2 Polyclonal Antibody, Unconjugated(bs-11331R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control (Black line): Molt4 (Black). Primary Antibody (green line): Rabbit Anti-TBR2antibody (bs-11331R) Dilution: 1µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody (white blue line): Goat antirabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

[IF=8.2] Wang Ziyi. et al. Targeting Myeloid Trem2 Reprograms the Immunosuppressive Niche and Potentiates

Checkpoint Immunotherapy in NASH-Driven Hepatocarcinogenesis. CANCER IMMUNOL RES. 2025 Aug;: FC; Mouse. 40748990

- [IF=6.208] Jia Wang. et al. FOXG1 Contributes Adult Hippocampal Neurogenesis in Mice. INT J MOL SCI. 2022 Jan;23(23):14979 IHC, IF; Mouse. 36499306
- [IF=4.3] Wang Yu-Xin. et al. PGC-1α Expands Neural Precursor Pool and Facilitates Cognitive Recovery Within AD Hippocampus Through the Regulation of Mitochondrial Dynamics. MOL NEUROBIOL. 2025 Jun;:1-23 IF,WB; Mouse. 40540175