

bs-11331R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

TBR2 Rabbit pAb**— DATASHEET —**

Host: Rabbit Clonality: Polyclonal GeneID: 8320 Target: TBR2 Immunogen: KLH conjugated synthetic peptide derived from human TBR2: 301-400/686. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: EOMES (eomesodermin homolog), also known as TBR2 (T-box-brain2), is the 686 amino acid human homolog of the mouse Eomes protein that contains one T-box DNA binding domain. Genes that contain T-box domains encode proteins that function as transcription factors and are often involved in the regulation of various developmental events. Localized to the nucleus and expressed in the developing brain, EOMES is thought to be involved in neuronal migration and division and may play a role in trophoblast development and gastrulation. Silencing of the EOMES gene can cause mutated or arrested development and may lead to microcephaly disorders which are characterized by reduced head circumference and a malformed brain.	Isotype: IgG SWISS: O95936	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test) Reactivity: Human, Rat (predicted: Mouse, Sheep, Cow, Chicken, Horse) Predicted MW.: 73 kDa Subcellular Location: Nucleus
--	---	---

— VALIDATION IMAGES —

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TBR2 Polyclonal Antibody, Unconjugated (bs-11331R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Blank control (Black line): Molt4 (Black). Primary Antibody (green line): Rabbit Anti-TBR2 antibody (bs-11331R) Dilution: 1µg / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF647 Dilution: 1µg / test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=6.208]** Jia Wang. et al. FOXG1 Contributes Adult Hippocampal Neurogenesis in Mice. INT J MOL SCI. 2022

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Jan;23(23):14979 IHC,IF ;Mouse. 36499306