bs-12785R

[Primary Antibody]

VTI1B Rabbit pAb




www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	lsotype: lgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500) IHC-F (1:100-500)
GenelD: 10490	SWISS: Q9UEU0	IF (1:100-500)
Target: VTI1B		Reactivity: Human, Mouse, Rat
Immunogen: KLH conjugated synthetic peptide derived from human VTI1B: 111-210/323.		(predicted: Pig, Cow)
Purification: affinity purified by	Protein A	
Concentration: 1mg/ml		Predicted MW.: ^{27 kDa}
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cell membrane ,Cytoplasm


— VALIDATION IMAGES

75 — 48[®] 48 — 48 — 5 25 _ - − ∨∏1 17 — 11 —

Sample: Hela Cell (Human) Lysate at 40 ug Primary: Anti- VTI1B (bs-12785R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 27 kD Observed band size: 27 kD



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (VT11B) Polyclonal Antibody, Unconjugated (bs-12785R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (VT11B) Polyclonal Antibody, Unconjugated (bs-12785R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.