bs-23029R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Dnmt3a Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 1788 SWISS: Q9Y6K1

Target: Dnmt3a

Immunogen: KLH conjugated synthetic peptide derived from human Dnmt3a:

51-150/912.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: CpG methylation is an epigenetic modification that is important for

embryonic development, imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA methylation is required for mammalian development. This gene encodes a DNA methyltransferase that is thought to function in de novo methylation, rather than maintenance methylation. The protein localizes to the cytoplasm and nucleus and its expression is

developmentally regulated. [provided by RefSeq, Mar 2016]

Applications: WB (1:500-2000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Mouse, Rat

(predicted: Human, Rabbit,


Sheep, Cow, Dog)

Predicted


100 kDa MW.:

Subcellular Location: Cytoplasm ,Nucleus


VALIDATION IMAGES



Sample: Lane 1: Ovary (Mouse) Lysate at 40 ug Lane 2: Ovary (Rat) Lysate at 40 ug Primary: Anti-Dnmt3a (bs-23029R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 120 kD Observed band size: 120 kD



Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DNMT-3 alpha) Polyclonal Antibody, Unconjugated (bs-23029R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DNMT-3 alpha) Polyclonal Antibody, Unconjugated (bs-23029R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

- SELECTED CITATIONS -

- [IF=6.8] Ruimeng Liu. et al. Protective role of curcumin on broiler liver by modulating aflatoxin B1-induced DNA methylation and CYPs expression. ECOTOX ENVIRON SAFE. 2023 Jul;260:115086 WB ;Chicken. 37269612
- [IF=3.699] Xiao-Tan Zhang . et al. Baicalin reversal of DNA hypermethylation-associated Klotho suppression ameliorates renal injury in type 1 diabetic mouse model. Cell Cycle. 2020;19(23):3329-3347 WB; Mouse. 33190590