

bsm-52099R**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Histone H2B Recombinant Rabbit mAb

— DATASHEET —

Host: Rabbit**Clonality:** Recombinant**GeneID:** 3018**Target:** Histone H2B**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].

Isotype: IgG**CloneNo.:** 3A6**SWISS:** P33778**Applications:** WB (1:500-2000)**IHC-P** (1:50-200)**IHC-F** (1:50-200)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**ICC/IF** (1:50)**Reactivity:** Human, Mouse, Rat


Predicted
MW.: 14 kDa

Subcellular
Location: Nucleus


— VALIDATION IMAGES —




Sample: Lane 1: Hela cell lysates Lane 2: PC-12 cell lysates Primary: Anti-Histone H2B (bsm-52099R) at 1/500 dilution Secondary: Goat Anti-Rabbit IgG - HRP at 1/5000 dilution
Predicted band size: 14 kD Observed band size: 14 kD




Paraformaldehyde-fixed, paraffin embedded (human liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Histone H2B) Monoclonal Antibody, Unconjugated (bsm-52099R) at 1:50 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Histone H2B) Monoclonal Antibody, Unconjugated (bsm-52099R) at 1:50 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



A431 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Histone H2B) monoclonal Antibody, Unconjugated



Blank control: Hela. Primary Antibody (green line): Rabbit Anti-Histone H2B antibody (bsm-52099R) Dilution: 1:50; Secondary Antibody: Goat anti-rabbit IgG-AF488 Dilution: 1:1000. Protocol The cells were fixed with 4%



Blank control: A431. Primary Antibody (green line): Rabbit Anti-Histone H2B antibody (bsm-52099R) Dilution: 1μg/10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody: Goat anti-rabbit IgG-AF488

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

(bsm-52099R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=16.6]** Daugird Timothy A. et al. Correlative single molecule lattice light sheet imaging reveals the dynamic relationship between nucleosomes and the local chromatin environment. NAT COMMUN. 2024 May;15(1):1-20 WB ;. 38755200
- **[IF=7.109]** Wu Yutong. et al. Reduced osteoclast-derived apoptotic bodies in bone marrow characterizes the pathological progression of osteoporosis. CELL DEATH DISCOV. 2023 Apr;9(1):1-9 WB ;Mouse. 37185334