

## IRF1 Recombinant Rabbit mAb

Catalog Number: bsm-52114R

Target Protein: IRF1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Recombinant

Clone No.: 3G7

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:50-200), IHC-F (1:50-200), IF (1:50-200), Flow-Cyt (1:50), ICC/IF (1:50)

Reactivity: Human, Mouse

Predicted MW: 37 kDa

Entrez Gene: 3659

Swiss Prot: P10914


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** IRF1 encodes interferon regulatory factor 1, a member of the interferon regulatory transcription factor (IRF) family. IRF1 serves as an activator of interferons alpha and beta transcription, and in mouse it has been shown to be required for double-stranded RNA induction of these genes. IRF1 also functions as a transcription activator of genes induced by interferons alpha, beta, and gamma. Further, IRF1 has been shown to play roles in regulating apoptosis and tumor-suppression.


### VALIDATION IMAGES




Sample: Lane 1: PC-12 cell lysates Lane 2: Jurkat cell lysates Primary: Anti-IRF1 (bsm-52114R) at 1/500 dilution Secondary: Goat Anti-Rabbit IgG - HRP at 1/5000 dilution Predicted band size: 37 kD Observed band size: 48 kD




Sample: Lane 1: Human THP-1 cell lysates Lane 2: Human Jurkat cell lysates Lane 3: Human MOLT4 cell lysates  
 Primary: Anti-IRF1 (bsm-52114R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
 Predicted band size: 37 kDa Observed band size: 46 kDa




Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IRF1) Monoclonal Antibody, Unconjugated (bsm-52114R) at 1:50 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IRF1) Monoclonal Antibody, Unconjugated (bsm-52114R) at 1:50 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IRF1) Monoclonal Antibody, Unconjugated (bsm-52114R) at 1:50 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: Jurkat. Primary Antibody (green line): Rabbit Anti-IRF1 antibody (bsm-52114R) Dilution: 1:50; Secondary Antibody : Goat anti-rabbit IgG-AF488 Dilution: 1:1000. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=4.36] Pengcheng Zhou. et al. Tiao-bu-fei-shen formula promotes downregulation of the caveolin 1-p38 mapk signaling pathway in COPD - Associated tracheobronchomalacia cell model. J ETHNOPHARMACOL. J Ethnopharmacol. 2022 Apr;115256 WB ; Mouse . 35367574

[IF=3.448] Yang T et al. Mechanism of berberine in treating Helicobacter pylori induced chronic atrophic gastritis through IRF8-IFN-γ signaling axis suppressing. Life Sci. 2020 Feb 22;248:117456. WB ; Rat . 32097666