

bs-4143R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Glutamine synthetase Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 2752	SWISS: P15104	IHC-F (1:100-500)
Target: Glutamine synthetase		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human Glutamine synthetase: 285-373/373.		ELISA (1:5000-10000)
Purification: affinity purified by Protein A		Reactivity: Mouse, Rat (predicted: Human, Pig, Sheep, Cow)
Concentration: 1mg/ml		Predicted MW.: 42 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cytoplasm
Background: Glutamine Synthetase catalyzes the conversion of ammonia and glutamate to glutamine. It is found in astrocytes as an octamer of identical 42 kDa subunits. The function of Glutamine Synthetase is the detoxification of brain ammonia. It also has an important role in the metabolic regulation of neurotransmitter glutamate. Because of the multiple functions and importance of Glutamine Synthetase in cellular metabolism, both catalytic activities and synthesis are highly regulated. The activity of Glutamine Synthetase is controlled by adenylation. Its activity is decreased in the cerebral cortex of brains affected by Alzheimer's disease, particularly in the vicinity of senile plaques. It is also decreased under conditions of glucose deprivation. The level of expression of Glutamine Synthetase is increased during ischemia in vivo or hypoxia in culture.		

— VALIDATION IMAGES —

Sample: Raji Cell (Human) Lysate at 40 ug
 Primary: Anti- Glutamine synthetase (bs-4143R)
 at 1/300 dilution Secondary: IRDye800CW Goat
 Anti-Rabbit IgG at 1/20000 dilution Predicted
 band size: 42 kD Observed band size: 42 kD



Tissue/cell: mouse brain tissue; 4%
 Paraformaldehyde-fixed and paraffin-
 embedded; Antigen retrieval: citrate buffer (
 0.01M, pH 6.0), Boiling bathing for 15min; Block
 endogenous peroxidase by 3% Hydrogen
 peroxide for 30min; Blocking buffer (normal goat
 serum, C-0005) at 37°C for 20 min; Incubation:
 Anti-Glutamine synthetase Polyclonal Antibody,
 Unconjugated(bs-4143R) 1:200, overnight at 4°C,
 followed by conjugation to the secondary
 antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=18.027]** Shikai Liu. et al. On-Demand Generation of Peroxynitrite from an Integrated Two-Dimensional System for Enhanced Tumor Therapy. ACS NANO. 2022;XXXX(XXX):XXX-XXX WB ;Human. 3566853
- **[IF=3.77]** Zhan Xie. et al. A novel model of subretinal edema induced by DL-alpha aminoadipic acid. EXP EYE RES. 2023

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Jan,;109388 WB ;Rabbit,Mouse,Rat. 36652968