

bsm-51153M**[Primary Antibody]****HGF Mouse mAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Mouse	Isotype: IgG	Applications: WB (1:800-1000)
Clonality: Monoclonal	CloneNo.: 9B6	Reactivity: Human
GeneID: 3082	SWISS: P14210	
Target: HGF		
Immunogen: KLH conjugated synthetic peptide derived from human HGF: 521-554/723.		Predicted MW.: 80 kDa
Purification: affinity purified by Protein G		Subcellular Location: Secreted
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: HGF is a pleiotropic protein in the plasminogen subfamily of S1 peptidases. HGF is a potent, mesenchymally-derived mitogen for mature parenchymal hepatocytes, and acts as a growth factor for a broad spectrum of tissues and cell types. HGF signals through a transmembrane tyrosine kinase receptor known as MET. Human HGF is secreted as an inactive 728 amino acid (aa) single chain propeptide. HGF regulates many physiological processes including proliferation, scattering, morphogenesis and survival. HGF is a crucial mitogen for liver regeneration processes, especially after partial hepatectomy and other liver injuries. This recombinant human HGF is produced by human cells. (Tag free) Biological activity: The activity was measured by its ability to stimulate the proliferation of the monkey epithelial cell line 4MBr-5. Reconstitution: Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile PBS containing 0.1% endotoxin-free recombinant human serum albumin.		

— VALIDATION IMAGES —

Sample: U937(Human) Cell Lysate at 30 ug
Primary: Anti- HGF (bsm-51153M) at 1/1000
dilution Secondary: IRDye800CW Goat Anti-
Mouse IgG at 1/20000 dilution Predicted band
size: 80 kD Observed band size: 80 kD