bs-9379R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn

techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 23609 **SWISS:** Q9H000

Target: RNF62/MKRN2

Immunogen: KLH conjugated synthetic peptide derived from human

MKRN2/RNF62: 261-360/416.

Purification: affinity purified by Protein A

RNF62/MKRN2 Rabbit pAb

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The Makorins are a family of putative ribonucleoproteins

containing two to four C3H zinc fingers that may confer RNA-binding. In addition, they contain a C3HC4 RING zinc finger that allows them to function as E3 ubiquitin ligases. Makorin-2, also known as RNF62, HSPC070 or MKRN2, is a widely expressed, evolutionarily conserved protein with four C3H-type zinc fingers (three at the N-terminus and one at the C-terminus), one RING-type zinc finger and a cysteine and histidine motif similar to that found in Makorin-1. In Xenopus, Makorin-2 functions, via PI 3-kinase/Akt signaling, as a negative regulator of neurogenesis. In humans, Makorin-2 is overexpressed in various cancer cell lines, suggesting a possible role of Makorin-2 in tumor progression. In addition, Makorin-2 is co-expressed with Raf-1 in the same tissues and cell

lines.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000)

Reactivity: (predicted: Human, Mouse,

Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW.: 47 kDa

Subcellular Nucleus Location: