bs-23650R

[Primary Antibody]

Vinculin Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 7414 SWISS: P18206

Target: Vinculin

Immunogen: KLH conjugated synthetic peptide derived from human Vinculin:

1031-1134/1134.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Vinculin is a cytoskeletal protein associated with cell-cell and cellmatrix junctions, where it is thought to function as one of several

interacting proteins involved in anchoring F-actin to the membrane. Defects in VCL are the cause of cardiomyopathy dilated type 1W. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Multiple alternatively spliced transcript variants have been found for this gene, but the biological validity of some variants has not been determined.

[provided by RefSeq, Jul 2008]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) ICC/IF (1:100-500)

Reactivity: Human, Mouse

(predicted: Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog,


Horse)

Predicted


MW.: 125 kDa

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse stomach tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Vinculin) Polyclonal Antibody, Unconjugated (bs-23650R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Vinculin) polyclonal Antibody, Unconjugated (bs-23650R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

— SELECTED CITATIONS –

• [IF=16.6] Feng Yuting, et al. Geometric constraint-triggered collagen expression mediates bacterial-host adhesion. NAT COMMUN. 2023 Dec;14(1):1-13 IF; Rat. 38071397