bs-23687R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Parkin Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 5071 **SWISS:** 060260

Target: Parkin

Immunogen: KLH conjugated synthetic peptide derived from human Parkin:

261-465/465.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Parkinson's Disease, the second most common neurodegenerative

disease after Alzheimer's Disease, is characterized by the loss of dopaminergic neurons and the presence of Lewy bodies (comprised of alpha synuclein and parkin inclusions). Autosomal Recessive Juvenile Parkinsonism (AR-JP) is a recently described form of Parkinson's Disease that has been linked to a gene that codes for parkin. Parkin, a 52 kDa protein, has a suggested role in the ubiquitin/proteasome pathway for protein degradation. The amino terminus bears sequence homology to ubiquitin while functionally it acts as a RING type ubiquitin protein ligase (E3) that coordinates the transfer of ubiquitin to substrate proteins, thus

targeting them for degradation by the proteasome.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human, Mouse

(predicted: Rat, Pig, Sheep,


Cow)

Predicted MW.: 51 kDa


Subcellular Cell membrane, Cytoplasm

Location: , Nucleus


VALIDATION IMAGES



Sample: Lane 1: Mouse Brain Lysates Lane 2: Mouse Cerebellum Lysates Lane 3: Mouse Hippocampus Lysates Primary: Anti-Parkin (bs-23687R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51kDa Observed band size: 51kDa



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Parkin) Polyclonal Antibody, Unconjugated (bs-23687R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Parkin) Polyclonal Antibody, Unconjugated (bs-23687R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

- SELECTED CITATIONS -

- [IF=18.2] Tingkui Zhao. et al. A Triple-Targeted Rutin-Based Self-Assembled Delivery Vector for Treating Ischemic Stroke by Vascular Normalization and Anti-Inflammation via ACE2/Ang1-7 Signaling. ACS CENTRAL SCI. 2023;XXXX(XXX):XXX-XXX WB;Rat. 37396868
- [IF=11.092] Juan Cen. et al. A Water-Soluble Quercetin Conjugate with Triple Targeting Exerts Neuron-Protective Effect

- on Cerebral Ischemia by Mitophagy Activation. ADV HEALTHC MATER. 2022 Sep;:2200817 WB,IHC;Rat, Human. 36071574
- [IF=9] An Xiaohong. et al. Inhibition of PDGFRβ alleviates endothelial cell apoptotic injury caused by DRP-1 overexpression and mitochondria fusion failure after mitophagy. CELL DEATH DIS. 2023 Nov;14(11):1-17 IF; MOUSE.
- [IF=5.75] Liu P. et al. Scalp Acupuncture Attenuates Brain Damage After Intracerebral Hemorrhage Through Enhanced Mitophagy and Reduced Apoptosis in Rats.. Front Aging Neurosci. 2021 Dec;13:718631-718631 IHC; Rat. 34987374
- [IF=6.055] Yun-Yi Zou. et al. Exercise intervention improves mitochondrial quality in non-alcoholic fatty liver disease zebrafish. FRONT ENDOCRINOL. 2023; 14: 1162485 WB; Zebrafish. 37284220