bs-19138R

[Primary Antibody]

ZFY19 Rabbit pAb




www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800	
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)	
Clonality: Polyclonal		IHC-P (1:100-500) IHC-F (1:100-500)	
GenelD: 84936	SWISS: Q96K21	IF (1:100-500)	
Target: ZFY19		Reactivity: Human, Mouse, Rat	
Immunogen: KLH conjugated synthetic peptide derived from human ZFY19: 101-200/471.		(predicted: Rabbit, Pig, Sheep, Cow, Dog, Horse)	
Purification: affinity purified by F	Protein A		
Concentration: 1mg/ml		Predicted MW.: ^{52 kDa}	
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated		Subcellular Location: Cytoplasm	


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

- VALIDATION IMAGES

Sample: MCF-7(Human) Cell Lysate at 30 ug Primary: Anti-ZFY19 (bs-19138R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kD Observed band size: 52 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFY19) Polyclonal Antibody, Unconjugated (bs-19138R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFY19) Polyclonal Antibody, Unconjugated (bs-19138R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.