bs-3327R

[Primary Antibody]

BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-PDK1 (Ser241) Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GenelD: 5170 **SWISS:** 015530

Target: PDK1 (Ser241)

Immunogen: KLH conjugated synthesised phosphopeptide derived from human

PDPK1 around the phosphorylation site of Ser241: AN(p-S)FV.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: PDK1 (3 Phosphoinositide Dependent Protein Kinase 1)

phosphorylates AGC kinases.

PDK1 activates conventional PKC and PKC zeta through phosphorylation of critical threonine residues in the activation loop. PDK1 also phosphorylates Protein Kinase B (PKB) at threonine 308 in the presence of phosphatidylinositol-3,4,5-trisphosphate. Active Akt inactivates Glycogen Synthase Kinase 3 (GSK3), eventually leading to the dephosphorylation and activation of glycogen synthase and the stimulation of glycogen synthesis. Because of the role that PDK plays in insulin-induced glycogen synthesis and PKC activation it is a potentially important target for metabolic drug research. There are three named

isoforms.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Rat (predicted: Human,


Mouse, Rabbit, Pig, Cow,

Chicken)

Predicted MW.: 61 kDa

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES



Tissue/cell: rat brain tissue; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-PDK1 Polyclonal Antibody, Unconjugated(bs-3327R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

- SELECTED CITATIONS -

• [IF=2.19] Lin, Yuan-xiang, et al. "Similar PDK1-AKT-mTOR pathway activation in balloon cells and dysmorphic neurons of typell focal cortical dysplasia with refractory epilepsy." Epilepsy Research (2015). IHC; Human. 25847349