

bs-9643R

[Primary Antibody]

C17orf64 Rabbit pAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Target: C17orf64

Immunogen: KLH conjugated synthetic peptide derived from human C17orf64: 32-130/236.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Applications: WB (1:500-2000)

IHC-P (1:100-500)


IHC-F (1:100-500)

IF (1:100-500)


Reactivity: Human, Rat
(predicted: Mouse, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW.: 27 kDa


— VALIDATION IMAGES —




Sample: Jurkat(Human) Cell Lysate at 30 ug
Primary: Anti- C17orf64 (bs-9643R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 27 kD Observed band size: 25 kD



Sample: U937 Cell (Human) Lysate at 30 ug
Primary: Anti- C17orf64 (bs-9643R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 27 kD Observed band size: 27 kD



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-C17orf64 Polyclonal Antibody, Unconjugated(bs-9643R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-C17orf64 Polyclonal Antibody, Unconjugated(bs-9643R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining