bs-0293G-RBITC

## [ Secondary Antibodies ]

## Goat Anti-Rat IgG H&L, RBITC conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Goat Clonality: Polyclonal	<b>Isotype:</b> IgG	Applications: IF (1:200-1000) Flow-Cyt (1:50-200) ICC/IF (1:100-1000)
Target: Goat Anti-Rat IgG H&L   Purification: affinity purified by Protein G   Concentration: 2.0 mg/ml   Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol.   Store at -20°C for one year. Avoid repeated freeze/thaw cycles.   Background: Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		Excitation Spectrum: 540nm Emission spectrum: 574nm <b>Reactivity:</b> Rat
functions; to bind to recruit other cells and variability of the IgG p	ponse. The IgG molecule has two sepa the pathogen that elicited the respons molecules to destroy the antigen. Th pool is generated by somatic recombin ecificities in an individual at a given ti be 1011 variants.	se and to ne nation