### bs-5809R

## [ Primary Antibody ]

# Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# Galectin 7 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 3963 **SWISS:** P47929

Target: Galectin 7

Immunogen: KLH conjugated synthetic peptide derived from human Galectin 7.:

31-100/136.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** The galectins are a family of beta-galactoside-binding proteins

implicated in modulating cell-cell and cell-matrix interactions. Differential and in situ hybridization studies indicate that this lectin is specifically expressed in keratinocytes and found mainly in stratified squamous epithelium. A duplicate copy of this gene (GeneID:653499) is found adjacent to, but on the opposite strand

on chromosome 19. [provided by RefSeq, Jul 2008].

Applications: ICC/IF (1:50)

Reactivity: Human (predicted: Mouse,


Rat)

Predicted MW.: 15 kDa

Subcellular Secreted ,Cytoplasm

Location: , Nucleus

#### VALIDATION IMAGES



HUVEC cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Galectin 7) polyclonal Antibody, Unconjugated (bs-5809R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.