bs-3810R

[Primary Antibody]

Villin Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 7429 **SWISS:** P09327

Target: Villin

Immunogen: KLH conjugated synthetic peptide derived from human Villin:

351-450/827.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Villin can cap, nucleate, sever and bundle actin in a calcium and phosphoinositide regulated manner. It is associated with the microvillar actin core bundle of intestinal and renal brush border implicated in adsorption. Villin is composed of six repeats, each containing 150 residues that together constitute the core domain followed by the carboxyl terminal headpiece domain of 87 residues. The core domain retains the calcium dependent capping nucleating and severing activity, whereas the headpiece domain contributes towards actin filament bundling and binding F actin, independently of Calcium. Function: Epithelial cell-specific Ca(2+)regulated actin-modifying protein that modulates the reorganization of microvillar actin filaments. Plays a role in the actin nucleation, actin filament bundle assembly, actin filament capping and severing. Binds phosphatidylinositol 4,5bisphosphate (PIP2) and lysophosphatidic acid (LPA); binds LPA with higher affinity than PIP2. Binding to LPA increases its phosphorylation by SRC and inhibits all actin-modifying activities. Binding to PIP2 inhibits actin-capping and -severing activities but enhances actin-bundling activity. Regulates the intestinal epithelial cell morphology, cell invasion, cell migration and apoptosis. Protects against apoptosis induced by dextran sodium sulfate (DSS) in the gastrointestinal epithelium. Appears to regulate cell death by maintaining mitochondrial integrity. Enhances hepatocyte growth factor (HGF)-induced epithelial cell motility, chemotaxis and wound repair. Upon S.flexneri cell infection, its actin-severing activity enhances actin-based motility of the bacteria and plays a role during the dissemination.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) **ELISA** (1:5000-10000)

Reactivity: Rat (predicted: Human,

Mouse, Pig, Cow, Chicken,

Dog, Horse)

Predicted 93 kDa

MW.:

Subcellular Cytoplasm

SELECTED CITATIONS —

- [IF=2.766] Wang et al. Characteristic and functional analysis of a newly established porcine small intestinal epithelial cell line. (2014) PLoS.One. 9:e110916 WB; Porcine. 25337908
- [IF=0] Cordeiro LA et al. In vitro interaction of spermatozoa with hen's sperm storage tubules. Journées du GdR 3606 Repro, Apr 2017, Tours, France. 75 p., 2017. (hal-01605736 IHC; hens (Gallus gallus). PRODINRA: 392709