[Primary Antibody]

phospho-Paxillin (Tyr31) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –			-9800	
Host: Rabbit	Isotype: IgG	Applications:		
Clonality: Polyclonal			HC-F (1:100-500) F (1:100-500)	
GenelD: 5829	SWISS: P49023		, , , , , , , , , , , , , , , , , , ,	
Target: Paxillin (Tyr31)			Reactivity: Mouse (predicted: Human, Rat, Sheep, Cow, Chicken, Horse)	
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human Paxillin around the phosphorylation site of Tyr31: TP(p-Y)SY.				
Purification: affinity purified by Protein A		Predicted	Predicted MW.: ^{64 kDa}	
Concentration: 1mg/ml		MW.: °		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular _C Location: ^C	ell membrane ,Cytoplasm	
Background: Paxillin is a 64 kDa cytoskeletal adapter protein involved in organisation and function of focal adhesions, which are critical to cell adhesion and migration. This in turn plays a role in a wide variety of processes including embryogenesis, organogenesis, wound repair, inflammation and cancer. Paxillin contains LD motifs, LIM domains, SH3 and SH2 binding domains that serve as docking sites for cytoskeletal proteins, tyrosine kinases (e.g., FAK, Pyk 2, Src), serine/threonine kinases, GTPase activating proteins and other adaptor proteins (e.g., Actin, Vinculin, Crk).		e as FAK,		

— VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse uterus); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-Paxillin (Tyr31)) Polyclonal Antibody, Unconjugated (bs-3744R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.